

České vysoké učení technické v Praze
Fakulta jaderná a fyzikálně inženýrská

Czech Technical University in Prague
Faculty of Nuclear Sciences and Physical Engineering

Ústav fyzikální chemie J. Heyrovského
Akademie věd České republiky

J. Heyrovský Institute of Physical Chemistry
Academy of Sciences of the Czech Republic

Doc. RNDr. Svatopluk Civiš, CSc

**ČASOVĚ ROZLIŠENÁ INFRAČERVENÁ EMISNÍ
SPEKTROSKOPIE S FOURIEROVOU TRANSFORMACÍ:
APLIKACE V PULZNÍCH VÝBOJÍCH A LASEROVÉ
ABLACI KOVŮ**

**TIME-RESOLVED FOURIER TRANSFORM INFRARED
EMISSION SPECTROSCOPY:
APPLICATION TO PULSED DISCHARGES AND LASER
ABLATION OF METALS**

Summary

Time-resolved Fourier transform spectroscopy (TR-FTS) is reviewed, with emphasis on synchronous FTS using continuously scanning interferometers. By using a high-resolution Bruker IFS 120 HR, a TR-FTS method has been developed with the help of a micro-controller Field Programmable Gate Array processor (FPGA), where a maximum of 64 time-resolved data are recorded with a preset time interval in a single scan of the interferometer. The time resolution is $1\ \mu\text{s}$, limited by the response time of the detector system used. This method has been applied to a pulsed discharge in a He and H_2 mixture to observe time profiles of H_3^+ and He_2 emission spectra.

Similar method of TR-FTS was extended for study of ablation products arising after irradiation of metal targets with a pulsed nanosecond ArF ($\lambda = 193\ \text{nm}$) laser. For data sampling so called $1/n$ interleaved sampling method was developed. A high-repetition rate laser (1.0 kHz) with a 15 mJ output power was focused on a rotating metal target inside a vacuum chamber (average pressure 10^{-3} Torr). The infrared atomic emission spectra of several metals (Au, Ag and Cu) have been observed in the $1200\text{--}4000\ \text{cm}^{-1}$ spectral region with a time profile showing maximum intensity in $10\ \mu\text{s}$ after a laser shot. Several new atomic transitions from Rydberg states has been observed and assigned for the first time. This yields precise energy values of a number of Au, Ag and Cu Rydberg levels some of which have not been reported previously.

Souhrn

Metoda časově rozlišené spektroskopie s Fourierovou transformací je popsána s důrazem na aplikace synchronní spektroskopie s Fourierovou transformací s použitím kontinuálně skenujícího interferometru. Pomocí metody časového rozlišení a pomocí mikroprocesoru FPGA společně s vysoce rozlišeným infračerveným interferometrem Bruker IFS 120 HR lze změřit až 64 časově rozlišených spekter (interferogramů) pro definované časové intervaly během jednoho celkového skenu interferometru. Maximální dosažené časové rozlišení je $1\ \mu\text{s}$, které je limitované rychlostí použitého detektoru (InSb). Tato metodika byla použita společně s pulzními výboji směs He, Ar, $(\text{CN})_2$ a H_2 . Vysoce rozlišená emisní spektra, včetně jejich časových průběhů byla získána pro některé vybrané radikály a molekulární ionty, např. $(\text{ArH}$, ArH^+ , H_3^+ , He_2 a CN).

Podobná technika časově rozlišené spektroskopie s Fourierovou transformací byla použita při studiu reakčních produktů vznikajících v ablační plazmě při ozařování kovového terče pulzním ArF ($\lambda = 193\ \text{nm}$) laserem. Pro sběr dat byla vyvinuta metodika prokládání $1/n$. Laser s vysokou opakovací frekvencí (1.0 kHz) a výstupním výkonem 15 mJ byl použit k ozařování rotujícího kovového terče umístěného uvnitř vakuové komory při tlaku 10^{-3} Torr. Byla změřena infračervená spektra atomárních přechodů řady kovů (Au, Ag a Cu) ve spektrálním rozsahu $1800\text{--}4000\ \text{cm}^{-1}$. Tímto způsobem byla identifikována celá řada zcela nových dosud neznámých a experimentálně dosud neidentifikovaných atomárních přechodů.

Klíčová slova

Emisní spektroskopie, časově rozlišená spektroskopie, FT spektroskopie, laserová ablace, infračervená emisní spektroskopie

Keywords

Emission spectroscopy, time resolved spectroscopy, Fourier transform spectroscopy, laser ablation, infrared emission spectroscopy

Contents

1. INTRODUCTION	6
1.1. Time-resolved Fourier transform infrared spectroscopy	6
1.2. Continuous scan systems: Synchronous triggering and data sampling . . .	7
2. CONTINUOUS SCAN SYSTEMS:	
APPLICATION WITH DISCHARGES	8
2.1. He discharge plasma	9
2.1.1 Introduction	9
2.1.2 Experimental	10
2.1.3 Observed spectra and analysis	10
2.2. Hydrogen containing discharge	15
2.2.1 Experimental	16
2.2.2 Results and discussion	16
2.3. (CN) ₂ -containing discharge	19
2.3.1 Observed CN spectra and their analysis	21
3. CONTINUOUS SCAN SYSTEMS: APPLICATION WITH LASERS	24
3.1. Interleaved sampling $1/n$	24
3.2. Synchronous triggering and data sampling	25
3.3. The experimental setup	27
4. TR FTIR LASER EMISSION SPECTROSCOPY	28
4.1. Introduction	28
4.2. Results for Au	29
4.3. Results for Ag	32
5. CONCLUSION	36
6. REFERENCES	37
7. CURRICULUM VITÆ	44

1. INTRODUCTION

1.1. Time-resolved Fourier transform infrared spectroscopy

Time-resolved spectroscopy (TRS) is a wide-spectrum technique used for studying the dynamics of chemical reactions, or the dynamic properties of molecules, radicals and ions in liquid, gas and solid states. In the infrared spectral range it can be achieved by using lasers [72], grating spectrometers [65] or by interferometers [50]. The presented report is focused on the development and application of a time resolved system based on commercially available continuously scanning high resolution interferometer and its modification for time resolved Fourier transform spectroscopy (TR-FTS) [42].

The main advantage of TR-FTS lies in obtaining spectra in wide wavenumber intervals. The speed of data acquisition is limited by the duration of the acquisition process and by the band width of the used detector.

There are basically two ways of obtaining the time-resolved spectra: the continuous scan and the non-continuous, step scan [50, 65, 72]. The continuous scan is best used when the duration of the observed phenomenon is longer than the time needed for carrying out one scan, *i. e.* for obtaining an interferogram up to the maximum trajectory difference (Rapid and ultrarapid scanning FT). Time-shifted individual scans provide a sequence of interferograms from which a conventional spectrum can be calculated. When using the rapid scanning and short distance mirror traversing, a time resolution from 1000 s to 1 ms can be reached.

A special approach to the time-resolved spectra of phenomena lasting from milliseconds to microseconds is the synchronous scanning FT technique [43]. This method, as well as the methods mentioned below, requires the possibility of initiating the reaction in a pulse mode, *e. g.* using a laser, electric discharge, electron bombardment, a UV discharge lamp, *etc.* [14]. The apparatus carries out a continuous scan and, during the pulse, it reads the signal from the detector corresponding to the position of the mirror and to the time from the beginning of the pulse reaction using the He-Ne laser fringe signals generated by the interferometer. This method is called stroboscopic interferometry [72]. After accumulating a sufficient amount of data and scans, the time-shifted interferograms are composed. The time mode is usually from 10 ms to 1 μ s.

A more favorable method of non-continuous scanning in steps (step-scan) is achieved in discrete jumps and the time-resolved data from each position can be recorded after each transient event [65]. Such a system is easy to couple with a pulse laser or a pulsed discharge. The step-scan spectrometers are commercially available and are used mainly for photolytic experiments in biology. The resolution of commercial step-scan type interferometers is limited to 0.1 cm^{-1} . A high resolution measurement with a step-scan type interferometer has been reported: a Connes type interferometer (CNRS Orsay) was used for the measurement of N_2 spectra with a resolution of 0.03 cm^{-1} [21].

1.2. Continuous scan systems: Synchronous triggering and data sampling

Continuously scanning spectrometers have been applied for time-resolved spectroscopy by several teams following the first report by Mantz [49]. Berg and Sloan [6] developed a compact data acquisition system for submicrosecond time-resolved FTS. Nakanaga Ito and Takeo [54] applied a pulse discharge system to a continuously scanning interferometer without any modification of the systems software. The pulsed discharge was triggered by a He–Ne laser fringe signal with an appropriate delay time. The system was applied to the measurement of the time profiles of a vibration-rotation absorption spectrum of discharged CO. Recently, Kawaguchi *et al.* [43] reviewed the methods of time-resolved Fourier transform infrared spectroscopy and its application to pulsed discharges and demonstrated the technique of FTS using a high-resolution Bruker IFS 120 HR supported by a microcontroller SX or Field Programmable Gate Array processor (FPGA) on He₂, ArH and ArH⁺ spectra. The same system was used for studying the products of ArF excimer laser ablation products [13, 15, 41].

The continuous scanning principle was the basis for data acquisition by a modified (Bruker IFS 120) spectrometer in our laboratory at the J. Heyrovský Institute of Physical Chemistry, and a similarly modified spectrometer was used in Okayama (Japan).

The data acquisition system can be described as follows:

The position of the traversing mirror of the Michelson interferometer is detected by reading the interference maxima of the He–Ne laser emission. The input signal in a cosine function shape is digitally processed into rectangular pulses and becomes the internal standard of the interferometer. The frequency of these rectangular pulses depends on the mirror speed. In the classic measurement mode, the frequency is usually 10 kHz with a pulse duration of 100 μ s. An external processor monitors the beginning of the He–Ne laser digital pulse, its order and the zero position of the mirror. During one pulse, the signal from the detector is read (30 or up to 64 readings), this is the so-called AD trigger [41]. These signals are shifted in time by Δt , where $\Delta t = 1$ or $2, 3 \dots \mu$ s.

In this way, a matrix $I(t_k, \delta_i)$ of intensity I in times t_k is acquired for the given optical path difference δ_i (i being the index of the selected optical path difference, from its zero to maximum values). A discharge pulse of variable length can be arbitrarily inserted into the data acquisition process (AD trigger). This process results in 30 to 64 reciprocally time-shifted interferograms.

Time resolved spectra are obtained by collecting data at various points between the zero-crossings and calculating the FT transformation for each such point. This system was utilized using a FPGA processor. The main role of the FPGA processor in our experiment was to create a discharge or laser pulse and AD trigger signals (the signal for data collection from the detector) synchronously with the He–Ne laser fringe signals from

the spectrometer (see Figure 1 and Figure 2). The FPGA processor also controls the data transmission from the digital input board to the PC.




Figure 1: A diagram of the time resolved Fourier transform spectrometer with FPGA microcontroller




Figure 2: Timing diagram for time resolved FT measurement. The scan and He-Ne fringe signals are supplied from Bruker 120 HR spectrometer. The velocity of the scanner is 10 kHz 100 μ s time intervals are produced. The discharge trigger is programmed using FPGA microcontroller. Maximum 64 interferograms (64 time shifted spectra) can be obtained during one scan.

2. CONTINUOUS SCAN SYSTEMS: APPLICATION WITH DISCHARGES

Figure 1 depicts the experimental arrangement used in presented study. Infrared emission was observed from a pulsed discharge of FT time-resolved measurements. The parent

compound hydrogen or $(\text{CN})_2$ was entrained in an inert carrier gas (He, Ar) and entered in the 20 cm long positive column discharge (or hollow cathode) tube with an inner diameter of 12 mm. The pulsed discharge was induced by a high voltage transistor switch HTS 81 (Behlke electronic GmbH, Frankfurt, Germany) between the stainless steel anode and grounded cathode. The plasma produced from the reaction mixture was cooled by flowing water in the outer jacket of the cell. The best conditions for the generation of radicals or ions were found to be $p(\text{He, Ar}) = 2\text{--}10$ Torr and 50 mTorr of parent molecules. The voltage drop across the discharge was 1000 V, with a pulse width of 20 or 40 μs and 0.5 A peak-to-peak current. The scanner velocity of the FT spectrometer was set to produce a 10 kHz He–Ne laser fringe frequency which was used to trigger the pulsed discharge. The recorded spectral range was 1800–4000 cm^{-1} with an optical filter, and a unapodized resolution of 0.07 or 0.025 cm^{-1} . The 32–100 scans were coadded so as to obtain a reasonable signal-to-noise ratio. The observed wavenumbers were calibrated using CO ground state rotation-vibration lines presenting in the spectra [31] as impurities.

2.1. He discharge plasma

2.1.1 Introduction

The He_2 molecule is known as the first Rydberg molecule, since its spectrum was reported in 1913. Many spectroscopic studies have been carried out as compiled in a book of Huber and Herzberg [37] and in the DiRef web site [7]. Most of the spectra of He_2 molecule have been observed in the visible and ultraviolet regions. Ginter and Ginter [26], Ginter et al. [27] compiled and analyzed the energy levels of Rydberg states originating from the electronic configurations $(1\sigma_g)^2(1\sigma_u)np\lambda(^3\Pi_g, ^3\Sigma_g^+)$ and $(1\sigma_g)^2(1\sigma_u)ns\sigma, nd\lambda(^3\Sigma_u^+, ^3\Sigma_u^+, ^3\Pi_u, ^3\Pi_u)$ by multichannel quantum defect theory, where n is the principal quantum number in the united atom molecular orbital designation. According to the energy levels listed in Refs. [26, 27], many electronic transitions are expected in the infrared region. However, observations of the infrared spectra so far have been limited to the three band systems below 8000 cm^{-1} :

- (1) $b^3\Pi_g\text{--}a^3\Sigma_u^+$ with the 0–0 band origin at 4750 cm^{-1} , studied by Hepner [32], Gloersen and Dieke [29], and [66],
- (2) $B^1\Pi_g\text{--}A^1\Sigma_u^+$ with the 0–0 band origin at 3501 cm^{-1} , studied by Solka et al. [73]
- (3) the $4f\text{--}3d$ band in 5100–5800 cm^{-1} spectral region, studied by Herzberg and Jungen [34].

The assignment of $4f\text{--}3d$ band was the first example concerning electronic states originating from the f -orbital electron.

The time-resolved Fourier transform spectroscopic system was applied for the observations of He₂ emission spectra produced by a pulsed discharge [36]. This method has enabled us to observe many electronic transitions in the infrared region, including the previously reported bands. The spectroscopic analysis of newly observed three bands and their time profiles are briefly reported.

2.1.2 Experimental

The spectra of He₂ were observed in emission from a hollow cathode discharge plasma. The hollow cathode stainless steel tube was 20 cm long with an inner diameter equal to 12 mm. The ac discharge was maintained by a high voltage transistor switch applied between the stainless steel anode and grounded cathode. The emission of He₂ has been also observed from a positive column, where lines from vibrationally excited states of $b^3\Pi_g$ were found to be more intense, compared with the case of the hollow cathode discharge. Here we report only the spectra obtained from the hollow cathode discharge, because of its higher efficiency in the production of the highly excited electronic states of He₂.

The plasma made from a pure helium was cooled down by flowing water or by liquid nitrogen in outer jacket of the cell. The best conditions for the generation of the He₂ were found to be $p(\text{He}) = 1.33 \text{ kPa}$ (10 Torr).

2.1.3 Observed spectra and analysis

Figure 3 shows a part of the observed time-resolved emission spectrum from a discharge in He. The discharge was initiated at time zero and turned off at 20 μs . For AD-converter triggers, we used 3 μsec for the zero offset and interval values, that is, AD conversion occurs every 3 μsec from the start of the discharge and all together 30 pulses cover 90 μsec . The strong line (5880 cm^{-1}) in Figure 3 belongs to the He atomic line ($4d-3p$) and is observed as two intense peaks. It may be noted that the second peak appears after the discharge is off, that is, it is due to the afterglow plasma. The other spectral lines in Figure 3 pertain to the $4f-3d$ transitions of He₂ which have been analyzed by Herzberg and Jungen [34].




Figure 3: A portion of the time-resolved spectrum observed by a pulse discharge in He with a pressure of 1.33 kPa (10 Torr). The discharge was applied in the interval of 0–20 μsec with a peak current of 0.5 A. The strongest peak belongs to atomic He line ($4d-3p$). Other lines pertain to $4d-3p$ transitions of He_2 .

Figure 4 shows an observed spectrum in the 2750–5600 cm^{-1} region, where we averaged all 30 spectra obtained by the time-resolved method. In the figure, the $b^3\Pi_g-a^3\Sigma_u^+$ $\nu = 0-0$ band is strongly observed in 4800 cm^{-1} region. Most of spectral lines in the 5200–5900 cm^{-1} region could be attributed to the $4f-3d$ band [73]. In the 3300 cm^{-1} region, the $B^1\Pi_g-A^1\Sigma_u^+$ $\nu = 0-0$ band found to be weak. From the time profile, it appears that the population in the singlet $B^1\Pi_g$ state decreased during the discharge period and increased in the afterglow, similarly to that observed for high-energy triplet states.




Figure 4: An observed spectrum of He_2 in the 2750–5600 cm^{-1} , where 30 time-resolved spectra (90 μsec) are averaged. The discharge condition is given in the caption of Figure 3

In addition to these already reported bands, some new bands were observed. In the 3200 cm^{-1} region, two series of lines have been observed with no Q -branch transitions. Rotational assignments are listed in Table 1 with the observed wavenumbers.

The analysis using the standard energy level expressions gave the rotational and centrifugal distortion constants, and the band origin (term energy) as listed in Table 2.

Spin splitting was not observed in these bands. The magnitude of the rotational constants was useful for identification of the electronic state. The band origin frequencies 3204.9 cm^{-1} and 3194 cm^{-1} of the two bands were consistent with those of the $g^3\Sigma_g^+ - d^3\Sigma_u^+$ ($\nu = 0-0$) and $h^3\Sigma_u^+ - g^3\Sigma_g^+$ ($\nu = 0-0$) transitions, respectively. Both the bands were identified for the first time in the infrared region, although electronic states involved have been observed by other electronic transitions in the visible region. Molecular constants determined in the previous studies are also listed in Table 2 for comparison.

Time profiles of observed spectral lines are depicted in Figure 5 for several bands. Except for the transitions from the $b^3\Pi$ state, the high-energy Rydberg states are produced strongly in the afterglow plasma. This means those states are more fragile during the discharge period and may not be observed strongly in a normal DC discharge.

N	$P(N)$	o.-c.	$R(N)$	o.-c.
$h^3\Sigma_u^+ - g^3\Sigma_g^+$				
0	3177.5569	0.0006		
2	3134.9969	-0.0005	3206.4133	0.0018
4	3107.2556	-0.0001	3235.5447	0.0000
6	3080.1473	-0.0010	3264.8626	-0.0064
8	3053.7916	0.0000	3294.3028	0.0089
10	3028.3000	0.0012	3323.7108	-0.0054
12	3003.7698	-0.0002	3353.0067	0.0013
14	2980.2783	-0.0002		
$g^3\Sigma_g^+ - d^3\Sigma_u^+$				
1	3190.4064	-0.0019	3232.9664	-0.0013
3	3160.7770	-0.0003	3259.9345	0.0005
5	3130.2548	0.0017	3285.6522	0.0016
7	3098.93068	0.0021	3130.0069	0.0006
9	3066.8917	-0.0002	3332.8839	-0.0017
11	3034.2243	-0.0031	3354.1697	0.0005
13	3001.0169	0.0015		

Table 1: Observed transitions of He_2 (cm^{-1}). N denotes the rotational quantum number neglecting spin in lower electronic states.




Figure 5: Observed time profiles of emission intensities of He_2

The pulsed discharge and multi-sampling system produce an interesting spectral fea-

	Present	Previous
<hr/>		
$(h^3\Sigma_u^+)$		(a)
B	7.14853(24)	7.149
$D \times 10^3$	0.5053(24)	0.574
$H \times 10^7$	-0.686(87)	
E		
<hr/>		
$(g^3\Sigma_g^+)$		(b)
B	7.096458(94)	7.0968(1)
$D \times 10^3$	0.53071(44)	0.538(7)
E	3204.8746(11)	
<hr/>		
$(d^3\Sigma_u^+)$		(c)
B	7.226364(88)	7.2286(15) ^a
$D \times 10^3$	0.51991(37)	0.532(3)
E	0.0	0.0
<hr/>		
(a) Huber and Herzberg [37]		
(b) Orth and Ginter [56]		
(c) Ginter [28]		

Table 2: Molecular constants of He_2 in the $h^3\Sigma_u^+$, $g^3\Sigma_g^+$ and $d^3\Sigma_u^+$ states (cm^{-1} units). Numbers in parentheses denote one standard deviation and applied to the last significant digits.

ture of He_2 in the infrared region. Especially, when the data sampling is carried out after turning off the discharge, intense emissions from many electronic bands are strongly observed. The analysis may provide information about high-energy Rydberg states including states originating from f -orbital electrons.

Figure 6 shows the energy level diagram of He_2 , where the energy values are represented relative to the $a^3\Sigma_u^+(v=0)$ state, which is located 144952 cm^{-1} (18 eV) above the repulsive ground $X^1\Sigma_g^+$ state. The observed transitions are demonstrated by arrows.




Figure 6: Energy level diagram of He_2 . The transitions observed in the present study are shown by arrows. The energy value is measured from the $a^3\Sigma_u^+(v=0)$ state; $n(>1)$ is the principal quantum number in a united atom molecular orbital designation. The ionization limit to He_2^+ is 34316 cm^{-1} .

2.2. Hydrogen containing discharge

Hydrogen and helium are the two most abundant elements in the universe. The hydrogen molecule and its various hydrides are the first source of aggregation and formation of interstellar matter. This process occurs in dense interstellar clouds, in star-forming regions and also in the atmosphere of some heavy planets (*e. g.*, Jupiter, Saturn, Uranus). The H_3^+ ion plays a dominant role in all these cases [20, 33, 51]. After the first laboratory spectroscopic detection of H_3^+ [55], a large number of laboratory studies have been published, of which about 20 were concerned with measuring new infrared spectra, describing about 800 transitions from a variety of vibrational bands in the spectral range between 1800 and 9000 cm^{-1} . It is apparent from the comprehensive evaluation and compilation study of H_3^+ spectroscopy by Lindsay and McCall [46] that most laboratory studies were carried out using absorption measurements. The only exception consists of the pioneering experiments of Majewski *et al.* [47, 48], who used a combination of a water-cooled, high-pressure, high-current emission hollow cathode together with an FT spectrometer. Majewski *et al.* used a high pressure of hydrogen gas (5–50 Torr) and high discharge current (up to 2.5 A) for production of H_3^+ . They obtained a very dense spectrum containing, in addition to H_3^+ , a large number of H atomic, H_2 valence, H_2 Rydberg, H_3 neutral, and also other unidentified transitions.

In this part we describe the application to the observation of H_3^+ , He and H emission produced by a pulsed discharge in a He/ H_2 mixture in the infrared spectral range. The use of time-resolved FT spectroscopy opens new pathways and new points of view in the study of the formation and decay processes inside the discharge plasma and permits description of the dynamics of the formation and decay of excited states of the H_3^+ ion [14].

2.2.1 Experimental

The emission spectra from a hollow cathode discharge plasma in He/H₂ mixtures were observed with the time-resolved Fourier transform high-resolution Bruker IFS 120 HR interferometer [14]. The hollow cathode stainless steel tube, covered with an outer glass jacket, was 25 cm long with an inner diameter of 12 mm. The He/H₂ plasma was cooled by liquid nitrogen in the outer jacket of the cell. The voltage drop across the discharge was 800 V, with a pulse width of 20 μ s and 0.6 A peak-to-peak current. The recorded spectral range was 1800–4000 cm⁻¹ with an optical filter, at an unapodized resolution of 0.1 cm⁻¹. Sixty-four scans were averaged to obtain a reasonable signal-to-noise ratio. The initial pressure of H₂ was 0.35 Torr and the He pressure was changed from 2 to 10.8 Torr.

The experiments were carried out with pulsed discharge with a width of 20 μ s. Because of the high pressures (up to 10 Torr) required for generation of H₃⁺ and thus, subsequently, the short relaxation times of the H₃⁺ ions, the measurement was carried out with maximum time resolution of 1 μ s, which is currently limited by the response time of a preamplifier of the InSb detector. Compared to previous measurements [36, 42], the data acquisition system was modified for enabling recording of 64 time-shifted spectra in a single scan.

2.2.2 Results and discussion

Helium has a higher ionization potential than H₂ (24.6 eV and 15.4 eV correspondingly; see Ref. [38]), so that the electron temperatures are higher when He predominates in the discharge. Because of the low proton affinity of He (1.9 eV) compared to H₂ (4.5 eV), the He buffer is chemically quite inert.

The low temperature emission spectra from a hollow cathode discharge in a He/H₂ mixture were found to contain only several of the low J and K transitions of H₃⁺, together with the atomic lines of He and H. No further lines of He⁺, H⁺ or molecular lines of H₂ or of neutral H₃ which also absorb in this area, were found [18, 76]. Figure 7 shows part of the observed time-resolved emission spectra of He (2129.83 cm⁻¹) and H (2148.79 cm⁻¹). Figure 7(b) depicts the H₃⁺ line $Q(1,0)$ at 2529.724 cm⁻¹, which belongs to $\nu_2 = 1 \rightarrow 0$ band.




Figure 7: Time-resolved infrared spectrum at 2125–2155 cm^{-1} (a) and 2528–2532 cm^{-1} (b), observed in a pulsed discharge in a He (10.8 Torr)/H₂ (0.4 Torr) mixture. The discharge was applied in a time interval of 0–20 μs with a peak current of 0.5 A.

Figure 8 shows time profiles of emission lines of He (a), H (a) and H₃⁺ (b). The absolute energy levels and Einstein coefficients A_{ij} for He, H and H₃ were taken from the NIST database [62] and from [40]. The vertical axis shows observed intensity divided by the Einstein coefficients A_{ij} and corresponds to the abundance in the upper state of the transition. It is noted that the emission from $n = 5$ of H was observed through the $n = 5-4$ transition, but other transitions from $n = 5$ are not observed because of the limited frequency range. Therefore, the abundance in the initial state of atomic transitions should be multiplied by a factor of 6 in the case of hydrogen. This sequence of processes was observed for all the lines in the spectrum; the time in which the maximum emissions were measured for the individual transitions of the same species (He, H₃⁺ and H) did not differ by more than $\pm 2 \mu\text{s}$.








Figure 8: Time-resolved emission profiles: (a) excited He $5f^1F^o-4d^1D$ transition at 2474.64 cm^{-1} (magnified by ten), (b) He $3p^3P^o-3s^3S$ transition at 2327.77 cm^{-1} (strongest line in the FTIR spectrum), (c) H_3^+ transition $Q(1,0)$ at 2529.72 cm^{-1} and (d) H Brackett α ($n = 5-4$) at 2467.60 cm^{-1} . The absolute intensity of emission was corrected by the corresponding Einstein coefficients A_{ij} .

Chemical processes in He/ H_2 plasma that can lead to the formation of H_3^+ are discussed extensively in literature (Eqns. 1–4; see Plašil et al. [58]):




In all our experiments no emission from H_3^+ has been detected in pure hydrogen discharge. Reaction (4) is exothermic to have enough energy to produce H_3^+ in $\nu_2 = 1$ state. However, since H_2 is known to be efficient for relaxing the vibrational excited state, the vibrationally excited H_3^+ in pure H_2 discharge will be relaxed by the collision with H_2 . It should be noted that the obtained radiative lifetime of H_3^+ is about 10 ms.

By adding a large amount of He the vibrational relaxation will be suppressed, and the H_3^+ emission becomes strong. At low helium pressures, H_3^+ is formed directly in the discharge through direct ionization of H_2 and subsequent processes (Eqns. 1,4). Formation and decay of excited H_3^+ occurs in μs time range. The calculated values for the deactivation rate $k_{ij} = (2..5) \times 10^5 \text{ s}^{-1}$ are several orders of magnitude higher than the Einstein coefficients A_{ij} for spontaneous emission of H_3^+ , and recombination process described later, which clearly demonstrates the efficiency of collision processes for deactivation of the H_3^+ excited states. In Figure 8, the second weaker maximum of the H_3^+ line was found at $45 \pm 2 \mu\text{s}$. This appears as a consequence of collisions of H_2 and metastable He with a long lived lifetime, as shown in Eqns. (2) and (4).

The entire process in the afterglow is concluded by formation of H, which could be observed on two lines (2148.7 and 2467.8 cm^{-1}) at times of $40\text{--}60 \mu\text{s}$ (Figure 8d). The two lines corresponding to transitions in excited atomic hydrogen were observed with two peaks (Figure 8d): a small peak at $12 \mu\text{s}$ and a high-intensity peak with a maximum at $39 \pm 1 \mu\text{s}$. The former peak may be explained by direct dissociation of H_2 through collision with electron. We expect that the second peak is due to hydrogen atom produced by recombination of H_3^+ with electron.

When the discharge is turned off, the electron energy becomes lower through collision, and the recombination rate will be increased. The recombination products of the reaction of H_3^+ with an electron are either three neutral hydrogen atoms or a hydrogen molecule together with a hydrogen atom. For the most important case of near-zero initial energy, *i. e.*, low-energy electrons and small internal excitation of the H_3^+ ion, the two reaction channels can be expected (Eqns. 5,6) [53, 74].



We observed two atomic H transitions in our spectra after electron recombination of H_3^+ (Figure 8); both originated from energy levels above 13 eV . This is more than the kinetic energy released in an energetically more advantageous process (Eq. 6). Because of the high pressure of He, excitation of H atoms can be expected in collisions with excited He atoms in the early afterglow.

2.3. (CN)₂-containing discharge

The CN free radical is observed in interstellar molecular clouds and the atmospheres of stars, planets and comets. It is also significant in numerous laboratory processes at high temperatures (flames, chemical reactions, discharges) where it is often formed from trace amounts of carbon and nitrogen. It is a very strong absorber/emitter of

radiation and its spectra, extending from the vacuum UV far into the infrared without significant gaps, provide a very useful tool for its detection and monitoring. A vast proportion of the available spectral data arises from the $A^2\Pi-X^2\Sigma^+$ and $B^2\Sigma^+-X^2\Sigma^+$ electronic transitions [60, 63] and the infrared transitions in the $X^2\Sigma^+$ ground electronic state [10, 35]. In our previous paper [35] we concentrated primarily on the measuring and analysis of $^{12}\text{C}^{14}\text{N}$ vibration-rotation bands for the sequences $\nu = (1-0)$ through $(9-8)$ which were observed in the spectral region 1800–2200 cm^{-1} with Fourier transform spectroscopy. From the point of view of the vibrational excitation, the most important information is obtained from vibronic data involving vibrational levels up to $\nu = 18$ [63]. Such high vibrational excitation corresponds to temperatures well above 45000 K thus indicating the potential use of CN in high temperature monitoring and the possibility of experimental determination of the molecular potential energy function [35].

Cerny et al. [10] analyzed fourteen vibronic bands of the $\Delta\nu = 1, 0, -1, -2$ spread out in the near infrared spectral range with $\nu = 0$ to 4 for $A^2\Pi$ electronic state. Kotlar et al. [44] carried out a perturbation analysis of data taken at the University of Berkeley, to give a deperturbed set of the constants for the $\nu = 0$ to $\nu = 12$ vibrational levels of the $A^2\Pi$ state. Prasad and Bernath [60] measured and analyzed the red system of CN by using a jet-cooled corona excited supersonic expansion in a spectral range of 16500–22760 cm^{-1} . They measured a total of 27 bands with $\nu = 8$ to 21 for $A^2\Pi$ electronic state. Furio et al. [24] used the laser fluorescence excitation spectra for the measurement of the $B^2\Sigma^+-A^2\Pi(\nu = 8, 7)$ band in the 20400 cm^{-1} spectral range and derived the constants for $\nu = 7$ of the $A^2\Pi$ state. Rehfuss et al. [64] used an FT spectrometer in the ultraviolet, visible and infrared region for a measurement of the CN spectrum. A total of 54 bands were observed throughout the red and infrared region from 16000 to 2500 cm^{-1} . The observed sequences include $\Delta\nu = +4, +3, +2, +1, 0, -1, -2$ and -3 with vibrational levels up to $\nu = 14$, where some sequences were not observed, due to small Franck–Condon factors and/or sensitivity of the spectrometer.

The 0–0 band of the $A^2\Pi-X^2\Sigma^+$ system appears at 9117 cm^{-1} . Since the vibrational frequency of CN is about 2042 cm^{-1} and 1813 cm^{-1} in the $X^2\Sigma^+$ and $A^2\Pi$ respectively, the $\Delta\nu = -1, -2$ and -3 sequences occur near 7000, 5000 and 3000 cm^{-1} respectively. In the region between 5000 and 2000 cm^{-1} , the vibronic transitions are rather unfavorable due to the Franck-Condon factors of 0.15–0.05 Prasad and Bernath [60], Sharp [70]. Furthermore one loses, compared with the 0–0 band, at least an additional factor of 20 due to the ν^3 dependence in the Einstein A coefficient. Thus a high resolution vibronic CN spectrum with a good signal-to-noise ratio for the $\nu = 3$ sequence band region has not been reported until now. Only a low resolution spectrum was weakly observed by Rehfuss et al. [64].

There is still a gap for the high resolution measurement and detailed analysis of the spectral bands concerning $\nu = 5$ to 6 of the $A^2\Pi$ state. The turning point in this measurement of CN in the infrared spectral range was the introduction of time resolved FT

spectroscopy [12]. This method makes it possible to distinguish the weak emission (or absorption) bands from strong bands appearing in the spectrum if the time profiles are different. In the case of CN, weak vibronic bands in the $5\ \mu\text{m}$ region were separated from strong long lived vibration-rotation bands. In this part a spectroscopic analysis of 7 newly observed $\Delta\nu = -3$ sequences bands: 0–3, 1–4, 2–5, 3–6, 4–7, 5–8 and 6–9 of the $A^2\Pi-X^2\Sigma^+$ transition is reported.

2.3.1 Observed CN spectra and their analysis

Figure 9 shows a part of the time-resolved FT spectra of emission from a discharge in a $(\text{CN})_2$ and He mixture, where the discharge pulse width was $20\ \mu\text{s}$. Thirty time-resolved spectra were obtained in one measurement with a time-interval of $3\ \mu\text{s}$, and 6 spectra are shown in Figure 9.




Figure 9: The time-resolved emission FT spectrum from a pulsed discharge in a $(\text{CN})_2$ and He mixture. The discharge pulse duration was $20\ \mu\text{s}$. The 30 time-resolved spectra were collected from $t = 0\text{--}90\ \mu\text{s}$ with a step of $3\ \mu\text{s}$. The spectra of C_2H_2 and C_2 were observed at 3300 and $3600\ \text{cm}^{-1}$.

The variation in intensity of the vibrational bands in the $X^2\Sigma^+$ state is low. On the other hand, relaxation of electronic transitions is as fast as expected from a short radiative lifetime. The wavenumber resolution of Figure 9 was $0.07\ \text{cm}^{-1}$, which was found to be insufficient for the analysis of the majority of the electronic transitions,

because in the band-head region, the lines remained unresolved and the fit of the spectra was unsatisfactory.

In another time-resolved measurement we used a 0.025 cm^{-1} resolution with a long discharge pulse ($40\text{ }\mu\text{s}$) in order to reach the maximum excitation and to set the system into a “steady state”. The data collection system was set using the offset time of $5\text{ }\mu\text{s}$ before the end of the pulse and the spectra were taken in $1\text{ }\mu\text{s}$ intervals. A series of experiments was carried out under this condition, while the basic parameters of the discharge, He pressure and discharge current were varied. A series of time-resolved FT spectra was measured in time intervals of $1\text{--}30\text{ }\mu\text{s}$, providing the time profile of CN relaxation from the $A^2\Pi$ state to the ground electronic state. The time-scale was short for the study of the relaxation of the vibration-rotation transitions in the ground electronic state, but is enough for observations of relaxation of atomic He, N, C lines, and the C_2 radical. From this vast complex spectra, the spectrum No. 15, as shown in Figure 10, was chosen for the present spectroscopic analysis, which was obtained $10\text{ }\mu\text{s}$ after the end of the discharge. Figure 11 shows typical time-profiles of a vibration-rotation transition in the $X^2\Sigma^+$ state and a vibronic transition of CN, together with the atomic lines. The lifetime of the $A^2\Pi\text{--}X^2\Sigma^+$ transition is an order of $10\text{ }\mu\text{s}$. However, the vibrational relaxation of CN in its ground state is significantly longer; even at a time of $25\text{ }\mu\text{s}$ after the discharge, the intensity of the vibrational fundamental band is still rising. The intensity of the He atomic line (2469.7 cm^{-1}) shows a fast decay, and lines of C and N atoms also relax with a speed comparable with atomic helium.




Figure 10: The emission spectrum of the CN radical in the spectral range 3–5 μm . The overall view of the CN $A^2\Pi-X^2\Sigma^+(\Delta\nu = -3)$ sequence with the 1–0 fundamental electronic vibration-rotation band present with a band origin at 2042 cm^{-1} and the other hot bands.

The rotational assignments of the $A^2\Pi-X^2\Sigma^+$: 0–3, 1–4, 2–5, 3–6, 4–7 bands were carried out according to the transition frequency calculations using molecular constants reported by Cerny et al. [10]. The $v = 5$ –8 and 6–9 bands were assigned by molecular constants from Kotlar et al. [44] who reported the Dunham parameters. The standard deviation of the fitting was 0.0009 cm^{-1} .

The time-resolved experiment itself was carried out in a wide range of time scales and with various discharge pulse lengths, thus enabling a complex study of the relaxation processes of the CN system in helium. Such a pulsed discharge gives a stronger emission for the $\Delta\nu = -3$ bands, compared with that of a DC discharge. In this study emission from the $\nu = 5$ and 6 vibrational levels of the $A^2\Pi$ state was observed for the first time. The observed intensity of each band was plotted against the energy value of the upper state of the transition to give a vibrational temperature of 6700 K for the $A^2\Pi$. The value does not change significantly after 10 μsec , because the vibrational relaxation is not fast. Similarly the vibrational temperature is found to be 6757 ± 534 K for the $X^2\Sigma^+$. Using the vibrational temperatures, we estimated the abundance of the $A^2\Pi$ to be 0.58% of that of the $X^2\Sigma^+$.




Figure 11: Time-profiles of the He(I) atomic line, P22 (10.5) line of the $A-X$, 5-8 band and R(16) line of the 1-0 fundamental vibrationrotation band, and nitrogen atomic line. The discharge pulse was 40 μs long. The time-resolved spectra were collected after 35 μs (5 μs before the end of the discharge pulse) with a step of 1 μs .

3. CONTINUOUS SCAN SYSTEMS: APPLICATION WITH LASERS

3.1. Interleaved sampling $1/n$

In the case of the measurement of time-resolved spectra in combination with a laser whose maximum repetition rate is slower than the interferometer mirror speed, there is no possibility of sampling at each individual trigger point of the He-Ne laser.

The lowest scanning speed of the interferometer is limited to the He-Ne laser fringe frequency of about 3 kHz. However, by utilizing the under-sampling condition, sampling several times more slowly becomes possible. Figure 12 shows the clock pattern for sampling in the present experiments where triggers for the pulse event and for the sampling are produced with a period of $1/n$ times the He-Ne laser fringe frequency. Complete interferograms are then obtained with n scans if the trigger point is changed for each scan. The time sequence shown in Figure 12 corresponds to the case of $n = 3$.

The maximum frequency of the used ArF laser is 1 kHz. The laser pulse is therefore repeated every 1000 μs . The minimum speed of the interferometer mirror is 3 kHz, then the digital signal produced by the He-Ne laser is repeated every 333.33 μs . In order to

obtain data in the maximum density, *i. e.* for every trajectory difference defined by the He–Ne laser, the complete record is taken during three scans (for the mirror speed 3 kHz and laser repetition frequency 1 kHz). The complete set of the time-resolved spectra (one complete interferogram) is acquired by three time-shifted scans.

3.2. Synchronous triggering and data sampling

FT data are taken at the zero crossing points of the He–Ne laser fringe signals, while the wavelength of the He–Ne laser is used for the measurement of path differences. The data are sampled at time intervals which correspond to a mirror movement of either one wavelength or half a wavelength, depending on the frequency range of the measurements (8000 or 16000 cm^{-1}). Time resolved spectra are obtained by collecting data at various points between the zero-crossings and calculating the FT transformation for each such point. This system was utilized using a Field Programmable Gate Array processor (FPGA). The main role of the FPGA processor in our experiment was to create a discharge or laser pulse and AD trigger signals (the signal for data collection from the detector) synchronously with the He–Ne laser fringe signals from the spectrometer. The FPGA processor also controls the data transmission from the digital input board to the PC.

Figure 12 depicts the timing chart (clock pattern) produced by the FPGA for the laser-pulse ablation method. The scan signal and He–Ne laser fringe signals are supplied by the Bruker 120 HR spectrometer and used as the system time standard. A discharge trigger pulse is produced at a width which is preset by the FPGA. AD triggers are also produced by the FPGA with a time offset value between the beginning of the laser pulse and the interval between pulses. In the present experiments we used a 60 μs offset and interval values covering, 30 μs when 30 AD triggers were supplied. A series of data signals corresponding to the AD triggers are stored and Fourier-transformed.




Figure 12: Timing diagram for the interleaved sampling. During the scan, the laser pulse and the AD trigger sampling are induced with a rate of $1/n$ times of the He–Ne laser fringe frequency. The complete interferograms are obtained after n scans ($n = 3$ here).

The maximum number of spectra taken by this method is 64. The time resolution is about $1 \mu\text{s}$, which is limited by the band width of the detector amplifier. The present system collects 64 times more data in comparison to the original Bruker system. This is possible because of rapid development in the field of PCs, their memory size and the writing speed of the hard disk. In the case of our current data collection program, we are able to store 64 interferograms in a single scan when the resolution is up to 0.03 cm^{-1} . For resolutions higher than 0.03 cm^{-1} , 30 time-resolved data are recorded simultaneously, which means the number of time-resolved data points can be varied according to the type of experiment and the memory capacity.

3.3. The experimental setup

The time resolution FTIR spectra were measured using the modified Bruker IFS 120 HR spectrometer (modified for the time-resolution scan of emission data) in a spectral range of 1800–6000 cm^{-1} using a CaF_2 beam splitter, and an InSb detector. The aperture size was 4 mm, the preamplifier gain was 3. The spectra were measured at a resolution of 0.1 cm^{-1} with a MID IR filter for the range of 1538–3500 cm^{-1} (number of scans from 1 to 10, zero filling 2, trapezoid apodization function).

The Bruker system was equipped with an analog-digital converter (ADC 4322: Analogic, USA), which was connected to a PC containing a programmable control processor of Field Programmable Gate Array – FPGA, (ACEX 1K: Altera, USA) set up at a frequency of 33 MHz, and digital input board PCI (2172C: Interface, Japan). The data collection process and synchronization with the laser were controlled by the FPGA processor programmed by QUARTUS II 7.1, Altera. The computer programs for data acquisition and fast FT transformation and displaying of the data were written in C++ language.

Time-resolved FTIR spectroscopy was applied for observations of the emission arising after the irradiation of metals with a pulsed nanosecond ArF ($\lambda = 193$ nm) laser. A high repetition rate ArF laser ExciStar S-Industrial V2.0 1000 (193 nm, laser pulse width 12 ns, frequency 1 kHz) with 15 mJ output power was focused on a rotating and linearly traversing gold target inside a vacuum chamber (average pressure 10^{-2} Torr). The infrared emission (axial distance from the target 10 mm) was focused into the spectrometer using a CaF_2 (100 mm) lens (see Figure 13). The emission was observed in the 1800–3500 cm^{-1} spectral region with a time profile showing maximum emission intensity at 9–11 μs after the laser shot.




Figure 13: Experimental set-up of the metal emission measurement.

For data sampling we used the so-called 1/3 sampling, where the scanner rate was set to produce a 3 kHz HeNe laser interference signal, the ArF laser oscillation was triggered, and 30 sets of time-resolved data were recorded with a preset time interval of $1\ \mu\text{s}$. Three scans were needed for a complete interferogram, and only 5 scans were coadded to improve the signal-to-noise ratio. The acquired spectra were post-zerofilled in the OPUS program and subsequently corrected by subtracting the blackbody background spectrum. The wavenumbers, line widths and their intensities were then obtained using the peak picking method (OPUS).

4. TR FTIR LASER EMISSION SPECTROSCOPY

4.1. Introduction

Pulsed laser ablation and depositing processes are currently frequently used techniques. Laser induced plasma at low fluence (typically $10\ \text{J}/\text{cm}^2$) has numerous applications, *e. g.* Pulsed Laser Deposition (PLD) or multi-elemental analysis. The latter technique, known as Laser Induced Plasma Spectroscopy (LIPS) or Laser-Induced Breakdown Spectroscopy (LIBS) consists of analyzing the light spectrum emitted from a plasma created on the sample surface by laser pulses. LIPS has many practical advantages over the conven-

tional methods of chemical analysis of elements and is consequently being considered for a growing number of applications [2, 3, 30, 45, 61].

Excimer lasers operating in near-ultraviolet regions with typical laser fluences of 1–30 J/cm² are used for many types of ablation [17]. The ablation plume arising after irradiation with fluences of nanosecond duration pulses is governed by a great number of very complex physical processes. During the laser pulse (with typical duration of 20 ns), the laser photons heat the sample and bring a part of its surface to the critical temperature. The heated material starts to boil explosively [52] and creates an emission plume consisting of ejected particles, atoms and ions. The particles inside the plume can themselves interact with the laser photons, which leads to a subsequent rise in the temperature of the ablation plume and to photochemical and photodissociation processes [68, p. 219].

The population of Rydberg states responsible for IR emission lines is governed mainly by collisional processes. The electrons created in the photodissociation processes can interact with the laser pulse via the electron-ion inverse bremsstrahlung, which again causes additional heating of the plume [75] and leads to the fast transition of the plume from ionized gas to plasma. The electrons escaping from the corona region cause a separation of charges, thereby inducing the ionized part of the plasma to accelerate. After the end of the laser pulse, the plume expands adiabatically. The electron-ion collision inside the plume can create excited ions. The electron-ion collision in the presence of a third body can result in their recombination leading to formation of atoms in highly excited Rydberg states [16]. A radiative cascade of these Rydberg states is then observed as the optical emission of the ablation plume.

The investigation of such emission is complicated by nonequilibrium and nonstationary conditions of the plasma for the excited states [1], so the information on population dynamics is only scarcely available for these states [67]. As an example of such data we report temporal evolution dynamics for each IR atomic line of the recorded spectra of metal atoms.

The properties of the observed plumes obtained by the ablation of different materials can eventually reflect the superposition of the ensemble processes described above. Here we report some results of a study focused on time-resolved spectra arising from 193 nm pulsed laser ablation of metallic (Au, Ag and Cu) targets in a 10⁻³ Torr vacuum. The atomic metal spectra were measured by a high resolution Fourier Transform infrared spectrometer specially modified for time-resolved measurements [13, 15].

4.2. Results for Au

The observed IR emission spectra of the Au atom are presented in Figure 14 at 10 μ s after the laser shot, when the time profile of the emission intensity is maximum for all the observed lines.




Figure 14: Some parts of the observed IR emission spectra of Au. The 2743.358 cm^{-1} and 2747.567 cm^{-1} values are given to the centers of gravity of the hyperfine patterns clearly seen in the second graph as double peaks.

Although the Au spectrum has been studied in various spectral domains for several decades [9, 19, 22, 23, 25, 39, 59], to our knowledge only one experimental study [25] concerning the studied $3\text{--}5\text{ }\mu\text{m}$ IR range is reported. As compared to George et al. [25] we observed several strong new Au lines in the $1800\text{--}4000\text{ cm}^{-1}$ domain. The most prominent IR lines observed for Au are listed in Table 3. Their half-widths at half-maxima (HWHM) are calculated from fitting to the Lorentzian shape.

Wavenumber (cm ⁻¹)	Intensity (arb.u)	HWHM (cm ⁻¹)	Decay time (μs)	Identification
2156.484	12679	0.098	5.24 ± 1.7*	8f _{7/2} → 8d _{5/2}
2193.030	38690	0.12	6.56 ± 0.61	8s _{1/2} → 8p _{1/2}
2428.358	8024	0.39	6.36 ± 1.4*	12s _{1/2} → 9p _{3/2}
2474.954	53951	0.13	5.25 ± 0.21	7d _{5/2} → 6f _{7/2}
2512.219	36631	0.14	5.73 ± 0.25	9p _{1/2} → 10d _{3/2}
2518.489	121588	0.13	5.56 ± 0.22	7d _{3/2} → 6f _{5/2}
2520.684	16574	0.16	6.83 ± 0.58	8p _{3/2} → 8d _{5/2}
2522.683	91622	0.13	5.70 ± 0.36	5f _{7/2} → 8d _{5/2}
2743.370	8780	0.39	5.41 ± 1.2*	?
2744.380	41786	0.12	5.96 ± 0.68	8s _{1/2} → 8p _{3/2}
2747.567	10249	0.13	8.52 ± 2.6*	12s _{1/2} → 9p _{1/2}
2749.6	6453	0.27	5.51 ± 1.8*	23p _{1/2,3/2} → 9d _{3/2}
3187.811	13199	0.093	6.03 ± 0.91*	(5d ⁹ 6s)6p _{3/2} → 6d _{5/2}
3828.96	9540	0.14	5.47 ± 2.0*	7f _{7/2} → 7d _{5/2}
3862.41	9370	0.13	2.98 ± 1.1*	9d _{3/2} → 5f _{5/2}
3866.606	7738	0.11	6.84 ± 0.83*	9d _{3/2} → 8p _{3/2}

Table 3: Experimental Au lines and their identification. The decay time, τ , was calculated by exponential fitting of the measured time profiles of the corresponding lines. Time profile of the lines denoted by asterisk demonstrates significant deviation from the exponential decay; τ values are roughly approximate

The decay time, τ given in the Table 3 was calculated by fitting of the measured time profiles of the corresponding lines. These profiles are given in Figure 15. The time decay of most of the strong lines is well described by exponential fitting, excepting the 2156.484 cm⁻¹ line which demonstrates a non-constant decay rate during the 30 μs after the laser shot. Some weaker lines demonstrate such behavior more clearly, their decay is not exponential (and is even non-monotonic), so their τ values are estimated in Table 3 in a rough approximation. Such a non-monotonic decay can be due to more complex population kinetics of the atomic Au states in the ablation plasma. We consider most of the observed lines to be due to transitions between the Rydberg $n = 5..10$ states of the valence electron outside the closed-shell 5d¹⁰ core.




Figure 15: The time profiles of some observed lines (dots) and their fit with exponential decay (solid curves)

4.3. Results for Ag

Some parts of the observed IR emission spectra of the Ag atom are presented in Figure 16 at $11\ \mu\text{s}$ after the laser shot, when the time profile of the emission intensity is maximum for all the observed lines. The most prominent IR lines observed for Ag are listed in Table 4. Their full widths at half-maxima (FWHM) are calculated from fitting to a Voigt profile, but under our conditions this profile does not differ much from the Lorentzian shape (see Ref. [13]).




Figure 16: A section of the the observed IR emission spectra of Ag.

Wavenumber (cm ⁻¹)	Intensity (arb. units)	SNR	FWHM (cm ⁻¹)	Decay time (μs)	Identification
1345.570(6)	14889	14.	0.125(55)	4.17(104)(b)	$(4d^{10})7d_{\frac{3}{2}} \leftarrow (4d^{10})6f_{\frac{5}{2}}$
1363.326(14)	3910	4.3	0.146(150)	5.90(609)(b)	$(4d^{10})5f \leftarrow (4d^{10})6g$
1460.115(6)	4530	5.6	0.111(035)	3.34(170)(b)	$(4d^{10})7p_{\frac{3}{2}} \leftarrow (4d^{10})8s_{\frac{1}{2}}$
2149.320(2)	6956	2.4	0.025(19)	7.17(206)(b)	$(4d^{10})7d_{\frac{5}{2}} \leftarrow (4d^{10})7f_{\frac{7}{2}}$
2153.988(4)	32525	9.4	0.031(21)	3.07(15)	$(4d^{10})7s_{\frac{1}{2}} \leftarrow (4d^{10})7p_{\frac{1}{2}}$
2154.599(2)	5958	2.2	0.029(11)	12.8(78)(b)	$(4d^{10})7d_{\frac{3}{2}} \leftarrow (4d^{10})7f_{\frac{5}{2}}$
2234.168(1)	74835	13.	0.035(6)	3.63(34)	$(4d^{10})7s_{\frac{1}{2}} \leftarrow (4d^{10})7p_{\frac{3}{2}}$
2417.757(2)	4880	5.9	0.030(14)	5.03(214)(b)	$(4d^{10})6d_{\frac{3}{2}} \leftarrow (4d^{10})8p_{\frac{1}{2}}$
2447.045(2)	5463	6.6	0.026(24)	4.88(81)(b)	$(4d^{10})6d_{\frac{5}{2}} \leftarrow (4d^{10})8p_{\frac{3}{2}}$
2477.833(1)	113278	10.	0.044(4)	4.27(29)	$(4d^{10})6d_{\frac{5}{2}} \leftarrow (4d^{10})5f_{\frac{7}{2}}$
2488.1560(9)	78947	7.0	0.048(4)	4.20(35)(a)	$(4d^{10})6d_{\frac{3}{2}} \leftarrow (4d^{10})5f_{\frac{5}{2}}$
2506.8196(5)	460526	41.	0.056(2)	3.87(20)	$(4d^{10})4f \leftarrow (4d^{10})5g$
2579.210(3)	5252	6.2	0.036(21)	5.14(202)(b)	$(4d^{10})7p_{\frac{3}{2}} \leftarrow (4d^{10})7d_{\frac{3}{2}}$
2584.3387(6)	48450	31.	0.041(3)	4.20(37)(b)	$(4d^{10})7p_{\frac{3}{2}} \leftarrow (4d^{10})7d_{\frac{5}{2}}$
2658.7911(9)	22693	14.	0.038(5)	4.15(39)(a)	$(4d^{10})7p_{\frac{1}{2}} \leftarrow (4d^{10})7d_{\frac{3}{2}}$
3386.152(1)	98370	9.2	0.059(6)	3.35(28)	$(4d^{10})6p_{\frac{3}{2}} \leftarrow (4d^{10})7s_{\frac{1}{2}}$
3589.552(5)	13825	15.	0.065(4)	3.15(11)(a)	$(4d^{10})6p_{\frac{1}{2}} \leftarrow (4d^{10})7s_{\frac{1}{2}}$

Table 4: Experimental Ag lines and their identification. The decay time, τ , was calculated by exponential fitting of the measured time profiles of the corresponding lines. The profiles denoted as ((a)) demonstrates significant deviation from the exponential decay; those denoted by (b) demonstrate the decay curves of essentially non-exponential form with a plateau or a second maximum; τ value is roughly approximate

We measured the emission spectrum at a different delay time, from 0 to 30 μs , after the laser shot. This allows us to measure the time profiles of the observed Ag lines. Some such profiles are shown in Figure 17. The temporal decay of some lines is well described by exponential fitting, while some lines display non-exponential (including some “plateaux” at 20–25 μs after the laser shot) and even non-monotonic behavior. Therefore their decay time, τ , values are estimated in Table 4 in a rough approximation; so, for essentially non-exponential decays, the standard deviation $\Delta\tau$ is of order of τ .

It should be noted that the decay times $\tau \simeq 1\text{--}10 \mu\text{s}$ given in Table 4 are not related to the radiative lifetimes of Ag atom levels which are at least two orders shorter [4, 5, 77]. The temporal dynamics shown in Figure 17 is due to a complex combination of the collisional cascade repopulation of the emitting levels [13] and the transfer processes in ablation products [41].




Figure 17: The time profiles of some observed Ag lines (dots) and their fit with exponential decay (solid curves)

The revised energy values E_i of some Ag terms are presented in Table 5. For the levels with $n \leq 6$ our values coincide with the reference values within the uncertainty intervals, but it is not the case for $n > 6$. However we consider our values preferable since they are extracted from spectra recorded with 0.02 cm^{-1} resolution while the reference values were obtained from spectra with resolution of $0.035\text{--}0.045 \text{ cm}^{-1}$ [57] and 0.06 cm^{-1} [8]

Term	Energy (cm ⁻¹)	Other sources
$(4d^{10})5f_{\frac{5}{2}}$	56691.275(2)	56691.4 [71], 56692.5 [69]
$(4d^{10})5f_{\frac{7}{2}}$	56691.397(4)	56691.4 [71], 56694.4 [69]
$(4d^{10})6p_{\frac{1}{2}}$	48297.402(2)	48297.402(3) [57]
$(4d^{10})6p_{\frac{3}{2}}$	48500.804(1)	48500.804(2) [57]
$(4d^{10})6d_{\frac{3}{2}}$	54203.119(2)	54203.119(2) [57]
$(4d^{10})6d_{\frac{5}{2}}$	54213.564(3)	54213.570(3) [57]
$(4d^{10})6f_{\frac{5}{2}}$	58045.481(7)	This work
$(4d^{10})6g$	58054.723(16)	This work
$(4d^{10})7s_{\frac{1}{2}}$	51886.954(1)	51886.971(2) [57]
$(4d^{10})7p_{\frac{1}{2}}$	54041.087(2)	54040.99(6) [8]
$(4d^{10})7p_{\frac{3}{2}}$	54121.059(2)	54121.129(5) [57]
$(4d^{10})8s_{\frac{1}{2}}$	55581.246(3)	55581.258(3) [57]
$(4d^{10})8p_{\frac{1}{2}}$	56620.876(3)	56620.72(6) [8]
$(4d^{10})8p_{\frac{3}{2}}$	56660.596(6)	56660.559(17) [57]
$(4d^{10})7d_{\frac{3}{2}}$	56699.911(2)	56699.768(3) [57]
$(4d^{10})7d_{\frac{5}{2}}$	56705.435(2)	56705.498(3) [57]
$(4d^{10})7f_{\frac{5}{2}}$	58854.510(3)	This work
$(4d^{10})7f_{\frac{7}{2}}$	58854.755(3)	This work

Table 5: Revised values of some levels of Ag I

5. CONCLUSION

The longstanding interests of our laboratory are the spectroscopic investigations of molecular ions, radicals or atoms which play a fundamental role in many plasma chemical processes, as well as in the reactions taking place inside of interstellar clouds or in stellar envelopes of giant stars of our universe.

The presented report is focused on the development and application of a time resolved system based on commercially available continuously scanning high resolution interferometer and its modification for time resolved Fourier transform spectroscopy.

The use of time-resolved FT spectroscopy opens new pathways and new points of view in study of the formation and decay processes inside the discharge or laser plasma.

Here, we are able to study the individual processes using atomic or molecular lines in a very wide spectral range of the high resolution FT technology, which has been simultaneously extended into the time dimension.

The only limitation is the sensitivity of the infrared FT technique, together with the considerable time required for the acquisition of the spectral data.

6. REFERENCES

- [1] C. Aragon and J. A. Aguilera. Characterization of laser induced plasmas by optical emission spectroscopy: A review of experiments and methods. *Spectrochim. Acta, Part B*, 63(9):893–916, Sept. 2008. ISSN 0584-8547.
- [2] D. Babánková, S. Civiš, and L. Juha. Chemical consequences of laser-induced breakdown in molecular gases. *Progress in Quantum Electronics*, 30(2-3):75 – 88, 2006. ISSN 0079-6727.
- [3] O. Barthélemy, J. Margot, M. Chaker, M. Sabsabi, F. Vidal, T. W. Johnston, S. Laville, and B. L. Drogoff. Influence of the laser parameters on the space and time characteristics of an aluminum laser-induced plasma. *Spectrochim. Acta, Part B*, 60(7-8): 905 – 914, 31 Aug. 2005. ISSN 0584-8547. 3rd International Conference on Laser Induced Plasma Spectroscopy and Applications (LIBS04), Torremolinos, SPAIN, SEP 28-OCT 01, 2004.
- [4] G. J. Bengtsson, P. Jönsson, J. Larsson, and S. Svanberg. Time-resolved spectroscopic studies of the $7p\ ^2P$ states of neutral silver following vuv excitation. *ZPD*, 22(1):437–439, 1991. ISSN 0178-7683.
- [5] J. Bengtsson, J. Larsson, and S. Svanberg. Hyperfine structure and radiative-lifetime determination for the $4d^{10}6p\ ^2P$ states of neutral silver using pulsed laser spectroscopy. *Phys. Rev. A*, 42(9):5457–5463, 1 Nov. 1990. ISSN 1050-2947.
- [6] P. A. Berg and J. J. Sloan. Compact standalone data acquisition system for sub-microsecond time-resolved fourier transform spectroscopy. *Rev. Sci. Instrum.*, 64(9): 2508–2514, Sept. 1993. ISSN 0034-6748.
- [7] P. F. Bernath and S. McLeod. DiRef, a database of references associated with the spectra of diatomic molecules. *J. Mol. Spectrosc.*, 207(2):287, JUN 2001. ISSN 0022-2852.
- [8] C. M. Brown and M. L. Ginter. Absorption spectrum of Ag I between 1540 and 1850 Å. *J. Opt. Soc. Am.*, 67(10):1323–1327, 1977. ISSN 0030-3941.
- [9] C. M. Brown and M. L. Ginter. Absorption spectrum of Au I between 1300 and 1900 Å. *J. Opt. Soc. Am.*, 68(2):243–246, 1978. ISSN 0030-3941.
- [10] D. Cerny, R. Bacis, G. Guelachvili, and F. Roux. Extensive analysis of the red system of the CN molecule with a high resolution fourier spectrometer. *J. Mol. Spectrosc.*, 73(1):154 – 167, 1978. ISSN 0022-2852.

- [11] J. Chalmers and P. Griffiths, editors. *Handbook of Vibrational Spectroscopy (5 Volume Set)*. Wiley, 1 edition, 3 2002. ISBN 9780471988472.
- [12] S. Civiš, T. Šedivcová-Uhliková, P. Kubelík, and K. Kawaguchi. Time-resolved fourier transform emission spectroscopy of $A^2\Pi-X^2\Sigma^+$ infrared transition of the CN radical. *Journal of Molecular Spectroscopy*, 250(1):20 – 26, July 2008. ISSN 0022-2852.
- [13] S. Civiš, I. Matulková, J. Cihelka, K. Kawaguchi, V. E. Chernov, and E. Y. Buslov. Time-resolved fourier-transform infrared emission spectroscopy of Au in the 1800–4000-cm⁻¹ region: Rydberg transitions. *Phys. Rev. A*, 81(1):012510, Jan. 2010. ISSN 1050-2947.
- [14] S. Civiš, P. Kubát, S. Nishida, and K. Kawaguchi. Time-resolved fourier transform infrared emission spectroscopy of H_3^+ molecular ion. *Chem. Phys. Lett.*, 418(4-6): 448–453, Feb.6 2006. ISSN 0009-2614.
- [15] S. Civiš, I. Matulková, J. Cihelka, P. Kubelík, K. Kawaguchi, and V. E. Chernov. Time-resolved fourier-transform infrared emission spectroscopy of Ag in the (1300–3600)-cm⁻¹ region: Transitions involving f and g states and oscillator strengths. *Phys. Rev. A*, 82(2):022502, Aug 2010. ISSN 1050-2947.
- [16] F. Claeysens, R. J. Lade, K. N. Rosser, and M. N. R. Ashfold. Investigations of the plume accompanying pulsed ultraviolet laser ablation of graphite in vacuum. *J. Appl. Phys.*, 89(1):697–709, 1 Jan. 2001. ISSN 0021-8979.
- [17] F. Claeysens, S. J. Henley, and M. N. R. Ashfold. Comparison of the ablation plumes arising from arf laser ablation of graphite, silicon, copper, and aluminum in vacuum. *J. Appl. Phys.*, 94(4):2203–2211, ‘5 Aug. 2003. ISSN 0021-8979.
- [18] P. B. Davies, M. A. Guest, and R. J. Stickland. Infrared laser spectroscopy of H_2 and D_2 Rydberg states. I. Application of the polarization model. *J. Chem. Phys.*, 93 (8):5408–5416, 15 Oct. 1990. ISSN 0021-9606.
- [19] G. J. Ding, R. C. Shang, L. F. Chang, K. L. Wen, Q. Hui, and D. Y. Chen. Experimental study of Au atom Rydberg states. *J. Phys. B*, 22(10):1555–1561, 28 May 1989. ISSN 0953-4075.
- [20] P. Drossart, J.-P. Maillard, J. Caldwell, S. J. Kim, J. K. G. Watson, W. A. Majewski, J. Tennyson, S. Miller, S. K. Atreya, J. T. Clarke, J. H. Waite, and R. Wagener. Detection of H_3^+ on Jupiter. *Nature*, 340(6234):539–541, AUG 17 1989. ISSN 0028-0836.

- [21] G. Durry and G. Guelachvili. N₂ (B–A) time-resolved fourier transform emission spectra from a pulsed microwave discharge. *J. Mol. Spectrosc.*, 168(1):82–91, Nov. 1994. ISSN 0022-2852.
- [22] S. F. Dyubko, V. A. Efremov, V. G. Gerasimov, and K. MacAdam. Millimetre-wave spectroscopy of Au I Rydberg states: S, P and D terms. *J. Phys. B*, 38(8):1107–1118, 28 Apr. 2005. ISSN 0953-4075.
- [23] J. C. Ehrhardt and S. P. Davis. Precision wavelengths and energy levels in gold. *J. Opt. Soc. Am.*, 61(10):1342–1349, 1971. ISSN 0030-3941.
- [24] N. Furio, A. Ali, P. J. Dagdigian, and H.-J. Werner. Laser excitation of the overlapping CN $B-A(8, 7)$ and $B-X(8, 11)$ bands: The relative phase of the $B-A$ and $B-X$ transition moments. *J. Mol. Spectrosc.*, 134(1):199 – 213, Mar. 1989. ISSN 0022-2852.
- [25] S. George, A. Grays, and R. Engleman, Jr. Spectrum of Au I in the infrared using a fourier-transform spectrometer. *J. Opt. Soc. Am. B*, 5(7):1500–1502, July 1988. ISSN 0740-3224.
- [26] D. S. Ginter and M. L. Ginter. Multichannel interactions in the $(1\sigma_g)^2(1\sigma_u)ns\sigma, nd\lambda$ ($^3\Sigma_u^+$, $^3\Sigma_u^+$, $^3\Pi_u$, $^3\Pi_u$) Rydberg structures of He₂. *J. Chem. Phys.*, 88(6):3761–3774, 15 Mar. 1988. ISSN 0021-9606.
- [27] D. S. Ginter, M. L. Ginter, and C. M. Brown. Multichannel interactions in the $(1\sigma_g)^2(1\sigma_u)np\lambda(^3\Pi_g, ^3\Sigma_g^+)$ Rydberg structures of He₂. *J. Chem. Phys.*, 81(12):6013–6025, 1984. ISSN 0021-9606.
- [28] M. L. Ginter. The spectrum and structure of the He₂ molecule: Part III. characterization of the triplet states associated with the UAO’s $3s$ and $2p\pi$. *J. Mol. Spectrosc.*, 18(3):321 – 343, 1965. ISSN 0022-2852.
- [29] P. Gloersen and G. H. Dieke. Molecular spectra of hydrogen and helium in the infrared. *J. Mol. Spectrosc.*, 16(1):191–204, 1965. ISSN 0022-2852.
- [30] A. Gomes, A. Aubreton, J. J. Gonzalez, and S. Vacqui. Experimental and theoretical study of the expansion of a metallic vapour plasma produced by laser. *J. Phys. D*, 37(5):689, 7 Mar. 2004. ISSN 0022-3727.
- [31] G. Guelachvili and K. R. Rao. *Handbook of Infrared Standards: With Spectral Maps and Transition Assignments Between 3 and 2600 μm (v. 1)*. Academic Press, 7 1986. ISBN 9780123053602.
- [32] L. Hepner, G.; Herman. Nouveau système de bandes d’émission de la molécule He₂ vers 4700 cm⁻¹. *C. R. Acad. Sci. Paris, Ser. B*, 243(20):1504–1506, 1956.

- [33] E. Herbst and W. Klemperer. The formation and depletion of molecules in dense interstellar clouds. *Astrophys. J.*, 185:505–534, Oct. 1973. ISSN 0004-637X.
- [34] G. Herzberg and C. Jungen. The $4f$ states of He_2 : A new spectrum of He_2 in the near infrared. *J. Chem. Phys.*, 84(3):1181–1192, 1 Feb. 1986. ISSN 0021-9606.
- [35] V. Horká, S. Civiš, V. Špirko, and K. Kawaguchi. The infrared spectrum of CN in its ground electronic state. *Collection of Czechoslovak Chemical Communications*, 69(1):73–89, 2004. ISSN 0010-0765.
- [36] Y. Hosaki, S. Civiš, and K. Kawaguchi. Time-resolved Fourier transform infrared emission spectroscopy of He_2 produced by a pulsed discharge. *Chem. Phys. Lett.*, 383(3-4):256–260, 8 Jan. 2004. ISSN 0009-2614.
- [37] K. Huber and G. Herzberg. *Molecular Spectra and Molecular Structure. IV. Constants of Diatomic Molecules*. Van Nostrand, 1 edition, 12 1979. ISBN 9780442233945.
- [38] W. T. Huntress, Jr. Laboratory studies of bimolecular reactions of positive ions in interstellar clouds, in comets, and in planetary atmospheres of reducing composition. *Astrophys. J., Suppl. Ser.*, 33(4):495–514, Apr. 1977. ISSN 0067-0049.
- [39] E. Jannitti, A. M. Cant, T. Grisendi, M. Pettini, and G. P. Tozzi. Absorption spectrum of Au I in the vacuum ultraviolet. *Phys. Scr.*, 20(2):156–162, 1979. ISSN 0281-1847.
- [40] L. Kao, T. Oka, S. Miller, and J. Tennyson. A table of astronomically important ro-vibrational transitions for the H_3^+ molecular ion. *Astrophys. J., Suppl. Ser.*, 77(2):317–329, Oct. 1991. ISSN 0067-0049.
- [41] K. Kawaguchi, N. Sanechika, Y. Nishimura, R. Fujimori, T. N. Oka, Y. Hirahara, A. Jaman, and S. Civiš. Time-resolved fourier transform infrared emission spectroscopy of laser ablation products. *Chem. Phys. Lett.*, 463(1–3):38–41, 22 Sept. 2008. ISSN 0009-2614.
- [42] K. Kawaguchi, O. Baskakov, Y. Hosaki, Y. Hama, and C. Kugimiya. Time-resolved Fourier transform spectroscopy of pulsed discharge products. *Chem. Phys. Lett.*, 369(3-4):293–298, 17 Feb. 2003. ISSN 0009-2614.
- [43] K. Kawaguchi, Y. Hama, and S. Nishida. Time-resolved Fourier transform infrared spectroscopy: Application to pulsed discharges. *J. Mol. Spectrosc.*, 232(1):1–13, July 2005. ISSN 0022-2852.
- [44] A. J. Kotlar, R. W. Field, J. I. Steinfeld, and J. A. Coxon. Analysis of perturbations in the $A^2\Pi-X^2\Sigma^+$ ”red” system of CN. *J. Mol. Spectrosc.*, 80(1):86 – 108, 1980. ISSN 0022-2852.

- [45] W.-B. Lee, J.-Y. Wu, Y.-I. Lee, and J. Sneddon. Recent applications of laser-induced breakdown spectrometry: A review of material approaches. *Appl. Spectrosc. Rev.*, 39(1):27–97, 2004. ISSN 0570-4928.
- [46] C. M. Lindsay and B. J. McCall. Comprehensive evaluation and compilation of H_3^+ spectroscopy. *J. Mol. Spectrosc.*, 210(1):60 – 83, Nov. 2001. ISSN 0022-2852.
- [47] W. A. Majewski, M. D. Marshall, A. R. W. McKellar, J. W. C. Johns, and J. K. G. Watson. Higher rotational lines in the ν_2 fundamental of the H_3^+ molecular ion. *J. Mol. Spectrosc.*, 122(2):341 – 355, Apr. 1987. ISSN 0022-2852.
- [48] W. A. Majewski, A. R. W. McKellar, D. Sadovskii, and J. K. G. Watson. New observations and analysis of the infrared vibrationrotation spectrum of H_3^+ . *Can. J. Phys.*, 72(11-12):1016–1027, Nov-Dec 1994. ISSN 0008-4204.
- [49] A. W. Mantz. Infrared multiplexed studies of transient species. *Appl. Spectrosc.*, 30(4):459–461, 1976. ISSN 0003-7028.
- [50] K. Masutani. *Time-resolved Mid-infrared Spectrometry Using an Asynchronous Fourier Transform Infrared Spectrometer*, pages 655–665. Volume 1 of Chalmers and Griffiths [11], 1 edition, 3 2002. ISBN 9780471988472.
- [51] B. J. McCall, T. R. Geballe, K. H. Hinkle, and T. Oka. Detection of H_3^+ in the Diffuse Interstellar Medium Toward Cygnus OB2 No. 12. *Science*, 279(5358):1910–1913, 20 Mar. 1998. ISSN 0036-8075.
- [52] A. Miotello and R. Kelly. Laser-induced phase explosion: new physical problems when a condensed phase approaches the thermodynamic critical temperature. *Appl. Phys. A*, 69(Suppl. S):S67–S73, Dec. 1999. ISSN 0947-8396. 5th International Conference on Laser Ablation COLA’99, GOTTINGEN, GERMANY, JUL 19-23, 1999.
- [53] O. Motret, J. M. Pouvesle, and J. Stevefelt. Spectroscopic study of the afterglow excited by intense electrical discharges in high-pressure helium hydrogen mixtures. *J. Chem. Phys.*, 83(3):1095–1100, 1985. ISSN 0021-9606.
- [54] T. Nakanaga, F. Ito, and H. Takeo. Time-resolved high-resolution ftir absorption spectroscopy in a pulsed discharge. *Chem. Phys. Lett.*, 206(1-4):73 – 76, 30 Apr. 1993. ISSN 0009-2614.
- [55] T. Oka. Observation of the infrared spectrum of H_3^+ . *Phys. Rev. Lett.*, 45(7):531–534, Aug. 1980. ISSN 0031-9007.
- [56] F. B. Orth and M. L. Ginter. The spectrum and structure of the He_2 molecule: Characterization of the triplet states associated with the UAO’s $4p\sigma$, $5p\sigma$, and $6p\sigma$. *J. Mol. Spectrosc.*, 61(2):282 – 288, 1976. ISSN 0022-2852.

- [57] J. C. Pickering and V. Zilio. New accurate data for the spectrum of neutral silver. *Eur. Phys. J. D*, 13(2):181–185, Feb. 2001. ISSN 1434-6060.
- [58] R. Plašil, J. Glošik, V. Poterya, P. Kudrna, J. Rusz, M. Tichý, and A. Pysanenko. Advanced integrated stationary afterglow method for experimental study of recombination of processes of H_3^+ and D_3^+ ions with electrons. *Int. J. Mass Spectrom.*, 218(2):105 – 130, 1 July 2002. ISSN 1387-3806.
- [59] J. R. Platt and R. A. Sawyer. New classifications in the spectra of Au I and Au II. *Phys. Rev.*, 60(12):866–876, Dec. 1941.
- [60] C. V. V. Prasad and P. F. Bernath. Fourier transform jet-emission spectroscopy of the $A^2\Pi_i-X^2\Sigma^+$ transition of CN. *J. Mol. Spectrosc.*, 156(2):327 – 340, Dec. 1992. ISSN 0022-2852.
- [61] L. J. Radziemski and D. A. Cremers, editors. *Lasers-induced Plasmas and Applications*. Marcel Dekker, New York, 1989. ISBN 0-8247-8078-7. Chap. 7, pp. 295–325.
- [62] Y. Ralchenko, A. Kramida, J. Reader, and N. Team. NIST atomic spectra database (version 3.1.5), 2008.
- [63] R. Ram, S. Davis, L. Wallace, R. Engleman, D. Appadoo, and P. Bernath. Fourier transform emission spectroscopy of the $B^2\Sigma^+-X^2\Sigma^+$ system of CN. *J. Mol. Spectrosc.*, 237(2):225 – 231, June 2006. ISSN 0022-2852.
- [64] B. D. Rehfuss, M.-H. Suh, T. A. Miller, and V. E. Bondybey. Fourier transform UV, visible, and infrared spectra of supersonically cooled CN radical. *J. Mol. Spectrosc.*, 151(2):437 – 458, Feb. 1992. ISSN 0022-2852.
- [65] C. Rödiger and F. Siebert. *Fast Time-resolved Mid-infrared Spectroscopy Using an Interferometer*, pages 625–640. Volume 1 of Chalmers and Griffiths [11], 1 edition, 3 2002. ISBN 9780471988472.
- [66] S. A. Rogers, C. R. Brazier, P. F. Bernath, and J. W. Brault. Fourier transform emission spectroscopy of the $b^3\Pi_g-a^3\Sigma_u^+$ transition of He_2 . *Mol. Phys.*, 63(5):901–908, 10 Apr. 1988. ISSN 0026-8976.
- [67] M. Rossa, C. A. Rinaldi, and J. C. Ferrero. Internal state populations and velocity distributions of monatomic species ejected after the 1064 nm laser irradiation of barium. *J. Appl. Phys.*, 105(6):063306, 15 Mar. 2009. ISSN 0021-8979.
- [68] H.-G. Rubahn. *Laser Applications in Surface Science and Technology*. Wiley, New York, 5 1999. ISBN 9780471984504.

- [69] U. I. Safronova, I. M. Savukov, M. S. Safronova, and W. R. Johnson. Third-order relativistic many-body calculations of energies and lifetimes of levels along the silver isoelectronic sequence. *Phys. Rev. A*, 68(6):062505, Dec. 2003. ISSN 1050-2947.
- [70] C. M. Sharp. The computation of Franck–Condon factors, r -centroids and associated quantities in the electronic transitions of diatomic molecules. *Astron. Astrophys. Suppl. Ser.*, 55:33–50, Jan. 1984. ISSN 0365-0138.
- [71] A. G. Shenstone. The arc spectrum of silver. *Phys. Rev.*, 57(10):894–898, May 1940.
- [72] G. D. Smith and R. A. Palmer. *Instrumental Aspects of Time-resolved Spectra Generated Using Step-scan Interferometers*, pages 641–654. Volume 1 of Chalmers and Griffiths [11], 1 edition, 3 2002. ISBN 9780471988472.
- [73] H. Solka, W. Zimmermann, A. Stahn, D. Reinert, and W. Urban. Observation of the $B^3\Pi_u-A^3\Sigma_g^+$ band of He_2 . *Mol. Phys.*, 60(5):1179–1182, 10 Apr. 1987. ISSN 0026-8976.
- [74] D. Strasser, L. Lammich, H. Kreckel, S. Krohn, M. Lange, A. Naaman, D. Schwalm, A. Wolf, and D. Zajfman. Breakup dynamics and the isotope effect in H_3^+ and D_3^+ dissociative recombination. *Phys. Rev. A*, 66(3):032719, Sept. 2002. ISSN 1050-2947.
- [75] A. Vertes, R. W. Dreyfus, and D. E. Platt. Modeling the thermal-to-plasma transitions for Cu photoablation. *IBM J. Res. Dev.*, 38(1):3–10, Jan. 1994. ISSN 0018-8646.
- [76] M. Vervloet and J. K. Watson. Improved infrared and visible emission spectra of the H_3 and D_3 molecules. *Journal of Molecular Spectroscopy*, 217(2):255 – 277, Feb. 2003. ISSN 0022-2852.
- [77] J. Zhankui, P. Jönsson, J. Larsson, and S. Svanberg. Studies on radiative lifetimes in the $4d^{10}ns\ ^2S$ and $4d^{10}nd\ ^2D$ sequences of neutral silver. *Z. Phys. D*, 17(1):1–14, 1990. ISSN 0178-7683.

7. CURRICULUM VITÆ

Doc. RNDr. Svatopluk Civiš, CSc

Born: May 14, 1955, České Budějovice

Education and qualification degrees:

High school education: Gymnasium (High school), České Budějovice, Czech Republic 1970–1974

Graduate Studies:

- Charles University, Prague Czech Republic 1974–1979
- Diploma Thesis in Chemistry 1979, Prague, Czech Rep.
- Doctoral Thesis in Chemistry 1986, Prague, Czech Rep.
- Associate Professor in Chemistry 2006, Prague, Charles University

Positions:

1980–1985: Faculty of Natural Sciences, Charles University, Prague — assistant researcher

1984–1989: Institute of Raw Materials, Kutná Hora

1989–1990: J. Liebig University Giessen, Germany, Junior Alexander von Humboldt Fellowship

1990–1991: J. Heyrovský Institute of Physical Chemistry — research scientist

1991–1992: Herzberg Institute of Astrophysics, NRC of Canada in Ottawa — research associate fellowship

1993–2005: J. Heyrovský Institute of Physical Chemistry, Prague — research scientist,

2005– J. Heyrovský Institute of Physical Chemistry — Head of Department of Spectroscopy

Svatopluk Civiš studied inorganic chemistry at the Charles University, Prague and obtained the diploma degree in 1979. He was an assistant researcher in a period of 1980 to 1985 at the Department of Inorganic chemistry of Charles University. In 1988 he was awarded an Alexander von Humboldt Fellowship at the J. Liebig University Giessen,

Germany. The aim of his fellowship was a laboratory study of molecular ions using a laser diode and a Doppler modulation technique.

In 1990–91 he worked at the J. Heyrovský Institute of Physical Chemistry of Czech Academy of Science, Prague. He built the first high resolution infrared diode laser spectrometer in former Czechoslovakia.

In 1991 he obtained a research associate fellowship at the Herzberg Institute of Astrophysics, NRC of Canada in Ottawa. His research at NRC was oriented in the field of new experimental methods of molecular spectroscopy. He applied a different infrared and microwave techniques (different frequency laser, diode laser, millimetre-wave, and infrared FT emission spectroscopy) for investigation of spectra of unstable species and ions of an astrophysical importance.

After two years in Canada (in 1993) he returned to the J. Heyrovský Institute of Physical Chemistry as a research officer and head of the spectroscopy and photochemistry group.

Since 2006 he is the head of the Department of Spectroscopy at J. Heyrovský Institute of Physical Chemistry, Prague, Czech Republic.

Invitations abroad:

- C.N.R., Istituto di Spettroscopia Molecolare and Università di Bologna, Dipartimento di Chimica G. Ciamician, Bologna, Italy, as visiting professor, *Microwave spectroscopy of protonated formaldehyde and its isotopomers*, (1994 — 3 months; 1995 — 2 months)
- C.N.R. at Université des Sciences et Technologies de Lille, centre d'Etudes et de Recherches Lasers et Applications, Laboratoire de Spectroscopie Hertzienne, Lille, France, visiting professor, *Microwave spectroscopy of negative ions*, (1995 — 5 months)
- Université des Sciences et Technologies de Lille, centre d'Etudes et de Recherches Lasers et Applications, Laboratoire de Spectroscopie Hertzienne, Lille, France, visiting professor, *Microwave spectroscopy of negative ions*, (1996 — 3 months; 1997 — 2 months)
- University of Fukuoka, Japan, visiting professor, *Fourier Transform spectroscopy of unstable species*, (1997 — 2 months)
- Université de Paris Sud, Laboratoire d'Infrarouge, Orsay Cédex, Paris, visiting professor, *Velocity modulation — Fourier Transform spectroscopy of unstable species*, (1997 — 1 month)

- Okayama university, Department of Chemistry, Faculty of Sciences, Japan, visiting professor, *Interstellar spectroscopy*, (2000 — 4 months).
- Okayama university, Department of Chemistry, Faculty of Sciences, Japan, visiting professor, *Laboratory and interstellar spectroscopy*, (2002 — 2 months).
- Okayama university, Department of Chemistry, Faculty of Sciences, Japan, VBL researcher, *Laboratory and interstellar spectroscopy*, (2003 — 2 months).
- Fukuoka university, Faculty of Sciences, Japan, JSPS visiting professor, *Laser diode spectroscopy of molecular complexes* (2009 — 1 month)

Main fields of research at J. Heyrovský Institute are:

- Infrared diode laser and Microwave spectroscopy of molecular ions and short lived species
- Environmental laser chemistry
- Chemistry and spectroscopy of atmosphere
- Laser research motivated by astrobiology problems
- Fourier transform spectroscopy of transient species
- Atomic spectroscopy

Grants and projects: S. Civiš participated in more than 20 national and international projects funded by various institutions.

Teaching and student training activities:

- 2002 – : Faculty of Natural Sciences, Charles University, Prague, Czech Republic
- 2005 – : Czech Technical University in Prague, Faculty of Nuclear Sciences and Physical Engineering, Prague, Czech Republic
- Supervisor of 9+2 diploma theses, 9 bachelor theses, 4+3 PhD theses

Publication activity:

- More than 20 invited talks and seminars presented abroad.
- More than 110 papers in refereed journals, approx. 623 citations, (413 without self-citation).